

CHICKLYTE® WS | HIDROLYTE® WS

Soluble Powder

Energetic-Electrolytic Balanced Association with Glycine and Probiotics





Each 100 g contains: Sodium chloride 11 000 mg, Potassium chloride 4000 mg, Sodium phosphate 5000 mg, Magnesium sulfate anhydrous 800 mg, Iron (as iron-AHM*) 334 mg, Manganese (as manganese - AHM*) 1576 mg, Copper (as copper-AHM*) 500 mg, Zinc (as zinc-AHM*) 431 mg, Dextrose monohydrate 56 000 mg, Dihydrate trisodium citrate 200 mg, Anhydrous citric acid 5000 mg, Glycine 8000 mg, Probiotics: 3 mg, Lactose monohydrate q.s. ad 100 g. *AHM: Methionine-Hydroxy Analogue.



Treatment and prevention of dehydration conditions, electrolytic unbalance and lack of energy caused by excessive loss due to diarrhea, vomiting, sweating, illness, high temperature, excessive breathing and urine. As preventive and support in diarrheas. Convalescence, post-vaccination, beak trimming and poultry reception, transport stress, heat stress, acidosis.

Dosage and Administration

Poultry: 1 kg/1000 L of water. Pigs: 1 kg/750 L of water. Other especies: Lactating calves and foals: Individual: 0.5 g/kg body weight at a rate of 30 - 40 g/L of water. Feedlot calves: 1 kg/500 L of water ad libitum. Lambs, kids and tuis: Individual: 10 g per ration of water. In group: 150 g/10 L of water ad libitum. Cattle, horses, rabbits and other minor animals: Individual according to weight at a rate of 40 g/L of drinking water.

Commercial Presentation

Sachet x 100 g, Sachet x 1 kg, Sachet x 15 g,

SENASA Registry

SENASA Perú: A.16.07.N.0249

